

Structure of R-salbutamol Sulfate

S. Pagola, P. Stephens (Physics & Astronomy, SUNY, Stony Brook,) and S. Cuffini (CEPROCOR, Córdoba, Argentina)

Abstract No. pago0430

Beamline(s): X3B1

Introduction: Salbutamol is a drug used in asthma therapy. The single enantiomer, R-salbutamol, has desirable bronchodilator, bronchoprotective and anti-edematous properties. However, racemic salbutamol can intensify allergic bronchospasm and promote eosinophil activation in asthmatic airways¹.

The crystal structure of the racemic mixture has been previously determined² by single crystal X-ray diffraction. We have characterized the crystal structure of R-salbutamol sulfate, using high resolution X-ray powder diffraction.

Methods and Materials: The X-ray powder diffraction pattern of a sample of R-salbutamol sulfate was collected at X3B1 beamline in a capillary. The pattern was indexed using the program ITO³, and a Le Bail fit was done using FULLPROF⁴. A strong candidate structure was found with our program PSSP⁵.

Results: The powder pattern was indexed to the monoclinic cell $a=28.084(3)$ Å, $b=6.1428(7)$ Å, $c=16.502(2)$ Å, $\alpha=90^\circ$, $\beta=95.534(7)^\circ$, $\gamma=90^\circ$, cell volume= $2833.5(5)$ Å³. The space group C 2 is consistent with the systematic absences observed. As the racemic mixture, the material has two inequivalent salbutamol molecules in the asymmetric unit and a sulfate ion. Each R-salbutamol molecule has 5 torsions. Thus, 28 structural parameters to locate the molecules and determine the conformations were calculated. This is substantially more complex than other structures solved at present using simulated annealing methods. Since the number of potential false minima increases rapidly with structural complexity, more work is required to verify that this is the correct solution.

After Rietveld refinement of a candidate structure, $R_{wp}=15.6\%$ and $R_I=6.8\%$. The structure is shown in Figure 1.

Acknowledgments: Research performed at the N.S.L.S., Brookhaven National Laboratory, which is supported by the US D.O.E., Division of Chemical Sciences and Division of Materials Sciences. The SUNY X3 beamline at the National Synchrotron Light Source is supported by the Division of Basic Energy Sciences of the US D.O.E. (DE-FG02-86ER45231).

References: ¹ C.P. Page and J. Morley, *J. Allergy Clin. Immunol.* **104**, p. 31-41, 1999. ² J. Par, M. Leger, M. Goursolle and M. Gadret, "Structure Cristalline du Sulfate de Salbutamol [tert-Butylamino-2(Hydroxy-4 hydroxymethyl-3-phenyl)-1 Ethanol. $\frac{1}{2}$ H₂O]", *Acta Cryst. B34*, p.1203-1208, 1978. ³ J.W. Visser, *J. Applied Cryst.* **2**, p. 89-95, 1969. ⁴ J. Rodríguez-Carvajal, *Abstracts of the Satellite Meeting on Powder Diffraction of the XV Congress of the IUCr*, Toulouse, France, 1990. ⁵ S. Pagola and P.W. Stephens, "Crystal Structure Solution of Organic Materials by the Simulated Annealing Method", *J. Applied Cryst.*, submitted for publication.




Figure 1. Representation of the structure of R-Salbutamol sulfate along [010].